Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims:

- 1. (Withdrawn) An isolated polypeptide selected from the group consisting of:
- a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-43,
- b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:22-23, SEQ ID NO:28, SEQ ID NO:30-32, SEQ ID NO:36-41 and SEQ ID NO:43,
- c) a polypeptide comprising a naturally occurring amino acid sequence at least 91% identical to the amino acid sequence of SEQ ID NO:5,
- d) a polypeptide comprising a naturally occurring amino acid sequence at least 93% identical to the amino acid sequence of SEQ ID NO:27,
- e) a polypeptide comprising a naturally occurring amino acid sequence at least 94% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:35 and SEQ ID NO:29,
- f) a polypeptide comprising a naturally occurring amino acid sequence at least 95% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:4, SEQ ID NO:11, and SEQ ID NO:20,
- g) a polypeptide comprising a naturally occurring amino acid sequence at least 96% identical to the amino acid sequence of SEQ ID NO:9 and SEQ ID NO:18,
- h) a polypeptide comprising a naturally occurring amino acid sequence at least 97% identical to the amino acid sequence selected from the group consisting of SEQ ID NO:26, SEQ ID NO:33, and SEQ ID NO:34,

- i) a polypeptide comprising a naturally occurring amino acid sequence at least 98% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:6 7,
- j) a polypeptide comprising a naturally occurring amino acid sequence at least 99% identical to the amino acid sequence of SEQ ID NO:16,
- k) a polypeptide consisting essentially of a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2-3, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12-15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, and SEQ ID NO:42,
- 1) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and
- m) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-43.
- 2. (Withdrawn) An isolated polypeptide of claim 1 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-43.
- 3. (Currently Amended) An isolated polynucleotide encoding a polypeptide, wherein the polypeptide consists of the amino acid sequence of SEQ ID NO: 13of claim 1.
- 4. (Currently Amended) An The isolated polynucleotide of claim 3 wherein the polynucleotide sequence consists of SEQ ID NO: 56 encoding a polypeptide of claim 2.
- 5. (Canceled).
- 6. (Currently Amended) A recombinant polynucleotide comprising a promoter sequence operably linked to a the polynucleotide of claim 3.
- (Currently Amended) A cell transformed with a the recombinant polynucleotide of claim
- 8.-10. (Canceled).

- 11. (Withdrawn) An isolated antibody which specifically binds to a polypeptide of claim 1.12.-13. (Canceled).
- 14. (Withdrawn currently amended) A method of detecting a target polynucleotide in a sample, said target polynucleotide <u>comprising having a sequence of a the polynucleotide of claim 3 claim 12</u>, the method comprising:
- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.
- 15. (Withdrawn) A method of claim 14, wherein the probe comprises at least 60 contiguous nucleotides.
- 16. (Withdrawn currently amended) A method of detecting a target polynucleotide in a sample, said target polynucleotide <u>comprising having a sequence of a the polynucleotide of claim</u> 3 claim 12, the method comprising:
- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.
- 17. (Withdrawn) A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.
- 18. (Withdrawn) A composition of claim 17, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-43.

- 19. (Withdrawn) A method for treating a disease or condition associated with decreased expression of functional KPP, comprising administering to a patient in need of such treatment the composition of claim 17.
- 20. (Withdrawn) A method of screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:
 - a) contacting a sample comprising a polypeptide of claim 1 with a compound, and
 - b) detecting agonist activity in the sample.

21.-22. (Canceled).

- 23. (Withdrawn) A method of screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:
 - a) contacting a sample comprising a polypeptide of claim 1 with a compound, and
 - b) detecting antagonist activity in the sample.

24.-25. (Canceled).

- 26. (Withdrawn) A method of screening for a compound that specifically binds to the polypeptide of claim 1, the method comprising:
- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

- 27. (Withdrawn) A method of screening for a compound that modulates the activity of the polypeptide of claim 1, the method comprising:
- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.
- 28. (Withdrawn currently amended) A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a the polynucleotide sequence of claim 3 claim 5, the method comprising:
- a) contacting a sample comprising the target polynucleotide with a compound, under conditions suitable for the expression of the target polynucleotide,
 - b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

- 29. (Withdrawn currently amended) A method of screening for potential toxicity of a test compound, the method comprising:
 - a) treating a biological sample containing nucleic acids with the test compound,
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a the polynucleotide of claim 3 claim 12 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a the polynucleotide of claim 3 claim 12 or fragment thereof,
 - c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample indicates potential toxicity of the test compound.
- 30. (Withdrawn) A method for a diagnostic test for a condition or disease associated with the expression of KPP in a biological sample, the method comprising:
- a) combining the biological sample with an antibody of claim 11, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex, and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.
- 31. (Withdrawn) The antibody of claim 11, wherein the antibody is:
 - a) a chimeric antibody,
 - b) a single chain antibody,
 - c) a Fab fragment,
 - d) a F(ab')2 fragment, or
 - e) a humanized antibody.

- 32. (Withdrawn) A composition comprising an antibody of claim 11 and an acceptable excipient.
- 33. (Canceled).
- 34. (Withdrawn) A composition of claim 32, further comprising a label.
- 35. (Canceled).
- 36. (Withdrawn) A method of preparing a polyclonal antibody with the specificity of the antibody of claim 11, the method comprising:
- a) immunizing an animal with a polypeptide consisting of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
 - b) isolating antibodies from the animal, and
- c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-43.

37.-43. (Canceled).

- 44. (Withdrawn) A method of detecting a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-43 in a sample, the method comprising:
- a) incubating the antibody of claim 11 with the sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-43 in the sample.

- 45. (Withdrawn) A method of purifying a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-43 from a sample, the method comprising:
- a) incubating the antibody of claim 11 with the sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) separating the antibody from the sample and obtaining the purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-43.
- 46. (Withdrawn) A microarray wherein at least one element of the microarray is a polynucleotide of claim 13.
- 47. (Withdrawn) A method of generating an expression profile of a sample which contains polynucleotides, the method comprising:
 - a) labeling the polynucleotides of the sample,
- b) contacting the elements of the microarray of claim 46 with the labeled polynucleotides of the sample under conditions suitable for the formation of a hybridization complex, and
 - c) quantifying the expression of the polynucleotides in the sample.
- 48. (Withdrawn) An array comprising different nucleotide molecules affixed in distinct physical locations on a solid substrate, wherein at least one of said nucleotide molecules comprises a first oligonucleotide or polynucleotide sequence specifically hybridizable with at least 30 contiguous nucleotides of a target polynucleotide, and wherein said target polynucleotide is a polynucleotide of claim 12.
- 49. (Withdrawn) An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 30 contiguous nucleotides of said target polynucleotide.

- 50. (Withdrawn) An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to at least 60 contiguous nucleotides of said target polynucleotide.
- 51. (Withdrawn) An array of claim 48, wherein said first oligonucleotide or polynucleotide sequence is completely complementary to said target polynucleotide.
- 52. (Withdrawn) An array of claim 48, which is a microarray.
- 53. (Withdrawn) An array of claim 48, further comprising said target polynucleotide hybridized to a nucleotide molecule comprising said first oligonucleotide or polynucleotide sequence.
- 54. (Withdrawn) An array of claim 48, wherein a linker joins at least one of said nucleotide molecules to said solid substrate.
- 55. (Withdrawn) An array of claim 48, wherein each distinct physical location on the substrate contains multiple nucleotide molecules, and the multiple nucleotide molecules at any single distinct physical location have the same sequence, and each distinct physical location on the substrate contains nucleotide molecules having a sequence which differs from the sequence of nucleotide molecules at another distinct physical location on the substrate.

56.-141.(Canceled).

- 142. (New) An isolated polynucleotide complementary to the polynucleotide of claim 3.
- 143. (New) An isolated polynucleotide complementary to the polynucleotide of claim 4.
- 144. (New) An RNA equivalent of the polynucleotide of claim 3.